

Local Equivalence of Sacksteder and Bourgain Hypersurfaces

Maks A. Akivis and Vladislav V. Goldberg

2000 *Mathematics Subject Classification*. Primary 53A20, Secondary 14M99.

Keywords and phrases. Gauss mapping, varieties with degenerate Gauss mappings, hypercubic, Sacksteder, Bourgain.

Abstract. Finding examples of tangentially degenerate submanifolds (submanifolds with degenerate Gauss mappings) in an Euclidean space R^4 that are noncylindrical and without singularities is an important problem of differential geometry. The first example of such a hypersurface was constructed by Sacksteder in 1960. In 1995 Wu published an example of a noncylindrical tangentially degenerate algebraic hypersurface in R^4 whose Gauss mapping is of rank 2 and which is also without singularities. This example was constructed (but not published) by Bourgain.

In this paper, the authors analyze Bourgain's example, prove that, as was the case for the Sacksteder hypersurface, singular points of the Bourgain hypersurface are located in the hyperplane at infinity of the space R^4 , and these two hypersurfaces are locally equivalent.

1. It is important to find examples of tangentially degenerate submanifolds in order to understand the theory of such manifolds. These examples prove the existence of tangentially degenerate submanifolds and help to illustrate the theory. The first known example of a tangentially degenerate hypersurface of rank 2 without singularities in R^4 was constructed by Sacksteder [S 60]. This example was examined from the differential geometry point of view by Akivis in [A 87]. In particular, Akivis proved that the Sacksteder hypersurface has no singularities since they “went to infinity”. In the same paper, Akivis presented a series of examples generalizing Sacksteder's example in R^4 , constructed a new series of examples of three-dimensional hypersurfaces $V^3 \subset P^n(\mathbb{R})$ of rank 2 whose focal surfaces are imaginary, and proved existence of hypersurfaces of this kind. Note that more examples of tangentially degenerate submanifolds without singularities can be found in [I 98, 99a, 99b]. The examples are essentially based on classical Cartan's hypersurfaces (see [Ca 39]).

Mori [M 94] claims that he constructed “a one-parameter family of complete nonruled deformable hypersurfaces in R^4 with rank $r = 2$ almost everywhere”. However, it follows immediately from his formulas that the hypersurfaces of his family are not only ruled hypersurfaces—they are cylinders.

Also much progress on the study of tangentially degenerate submanifolds over the complex numbers has been made in [GH 79], [L 99], and [AGL 00]. In these papers and in the papers [FW 95], [W 95], and [WZ 99] one can find more examples of tangentially degenerate submanifolds over the complex numbers.

Recently Wu [W 95] published an example of a noncylindrical tangentially degenerate algebraic hypersurface in an Euclidean space R^4 which has a degenerate Gauss mapping but does not have singularities. This example was constructed (but not published) by Bourgain (see also [I 98, 99a, 99b]). In

the present paper, we investigate Bourgain's example from the point of view of the paper [A 87] (see also Section 4.7 of our book [AG 93]). In particular, we prove that, as was the case for the Sacksteder hypersurface, the Bourgain hypersurface has no singularities since they "went to infinity". Namely this analysis suggested an idea that Bourgain's and Sacksteder's examples must be equivalent. Moreover, this analysis showed that a hypersurface constructed in these examples is torsal, i.e., it is stratified into a one-parameter family of plane pencils of straight lines.

In addition, at the end of our paper we prove that the examples of Bourgain and Sacksteder are locally equivalent.

2. In Cartesian coordinates x_1, x_2, x_3, x_4 of the Euclidean space R^4 , the equation of the Bourgain hypersurface B is

$$x_1 x_4^2 + x_2(x_4 - 1) + x_3(x_4 - 2) = 0 \quad (1)$$

(see [W 95] or [I 98, 99a, 99b]). Equation (1) can be written in the form

$$x_1 x_4^2 + (x_2 + x_3)x_4 - (x_2 + 2x_3) = 0. \quad (2)$$

Make in (2) the following admissible change of Cartesian coordinates:

$$x_2 + x_3 \rightarrow x_2, \quad x_2 + 2x_3 \rightarrow x_3.$$

Then equation (2) becomes

$$x_1 x_4^2 + x_2 x_4 - x_3 = 0. \quad (3)$$

Introduce homogeneous coordinates in R^4 by setting $x_i = \frac{z_i}{z_0}$, $i = 1, 2, 3, 4$. Then equation (3) takes the form

$$f = z_1 z_4^2 + z_0 z_2 z_4 - z_0^2 z_3 = 0. \quad (4)$$

Equation (4) defines a cubic hypersurface F in the space $\overline{R}^4 = R^4 \cup P_\infty^3$ which is an enlarged space R^4 , i.e., it is the space R^4 enlarged by the hyperplane at infinity P_∞^3 (whose equation is $z_0 = 0$).

Denote by A_α , $\alpha = 0, 1, 2, 3, 4$, fixed basis points of the space $\overline{R}^4$. Suppose that these points have constant normalizations, i.e., that $dA_\alpha = 0$. An arbitrary point $z \in \overline{R}^4$ can be written in the form $z = \sum_\alpha z_\alpha A_\alpha$. We will take a proper point of the space $\overline{R}^4$ as the point A_0 , and take points at infinity as the points A_1, A_2, A_3, A_4 .

Equation (4) shows that the proper straight line $A_0 \wedge A_4$ defined by the equations $z_1 = z_2 = z_3 = 0$ and the plane at infinity defined by the equations $z_0 = z_4 = 0$ belong to the hypersurface F defined by equation (4).

We write the equations of the hypersurface F in a parametric form. To this end, we set

$$z_0 = 1, \quad z_4 = p, \quad z_1 = u, \quad z_3 = pv.$$

Then it follows from (4) that

$$z_2 = v - pu.$$

This implies that an arbitrary point $z \in F$ can be written as

$$z = A_0 + uA_1 + vA_2 + p(A_4 - uA_2 + vA_3). \quad (5)$$

The parameters p , u , v are independent nonhomogeneous parameters on the hypersurface F .

3. Let us find singular points of the hypersurface F . Such points are defined by the equations $\frac{\partial f}{\partial z_\alpha} = 0$. It follows from (4) that

$$\begin{cases} \frac{\partial f}{\partial z_0} = z_2 z_4 - 2z_0 z_3, \\ \frac{\partial f}{\partial z_1} = z_4^2, \quad \frac{\partial f}{\partial z_2} = z_0 z_4, \quad \frac{\partial f}{\partial z_3} = -z_0^2, \\ \frac{\partial f}{\partial z_4} = 2z_1 z_4 + z_0 z_2. \end{cases} \quad (6)$$

All these derivatives vanish simultaneously if and only if $z_0 = z_4 = 0$. Thus the 2-plane at infinity $\sigma = A_1 \wedge A_2 \wedge A_3$ is the locus of singular points of the hypersurface F .

Consider a point $B_0 = A_0 + pA_4$ on the straight line $A_0 \wedge A_4$. By (4), to the point B_0 there corresponds the straight line $a(p)$ in the 2-plane at infinity σ , and the equation of this straight line is

$$p^2 z_1 + pz_2 - z_3 = 0. \quad (7)$$

The family of straight lines $a(p)$ depends of the parameter p , and its envelope is the conic C defined by the equation

$$z_2^2 + 4z_1 z_3 = 0. \quad (8)$$

The straight line $a(p)$ is tangent to the conic C at the point

$$B_1(p) = A_1 - 2pA_2 - p^2A_3. \quad (9)$$

Equation (9) is a parametric equation of the conic C . The point

$$\frac{dB_1}{dp} = -2(A_2 + pA_3) \quad (10)$$

belongs to the tangent line to the conic C at the point $B_1(p)$.

Consider the 2-planes $\tau = B_0 \wedge B_1 \wedge \frac{dB_1}{dp}$. Such 2-planes are completely determined by the location of the point B_0 on the straight line $A_0 \wedge A_4$, and

they form a one-parameter family. All these 2-planes belong to the hypersurface F . In fact, represent an arbitrary point z of the 2-plane τ in the form

$$\begin{aligned} z &= \alpha B_0 + \beta B_1 - \frac{1}{2}\gamma \frac{dB_1}{dp} \\ &= \alpha A_0 + \beta A_1 + (-2p\beta + \gamma)A_2 + (-p^2\beta + p\gamma)A_3 + p\alpha A_4. \end{aligned} \quad (11)$$

The coordinates of the point z are

$$z_0 = \alpha, \quad z_1 = \beta, \quad z_2 = \gamma - 2p\beta, \quad z_3 = p(\gamma - p\beta), \quad z_4 = p\alpha. \quad (12)$$

Substituting these values of the coordinates into equation (4), one can see that equation (4) is identically satisfied. Thus the hypersurface F is foliated into a one-parameter family of 2-planes $\tau(p) = B_0 \wedge B_1 \wedge \frac{dB_1}{dp}$.

In a 2-plane $\tau(p)$ consider a pencil of straight lines with center B_1 . The straight lines of this pencil are defined by the point B_1 and the point $B_2 = A_2 + pA_3 + q(A_0 + pA_4)$. The straight lines $B_1 \wedge B_2$ depend on two parameters p and q . These lines belong to the 2-plane $\tau(p)$, and along with this 2-plane they belong to the hypersurface F . Thus they form a foliation on the hypersurface F .

We prove that this foliation is a Monge-Ampère foliation. In the space $\overline{R}^4$, we introduce the moving frame formed by the points

$$\begin{cases} B_0 = A_0 + pA_4, \\ B_1 = A_1 - 2pA_2 - p^2A_3, \\ B_2 = A_2 + pA_3 + qA_0 + pqA_4, \\ B_3 = A_3, \\ B_4 = A_4. \end{cases} \quad (13)$$

It is easy to prove that these points are linearly independent, and the points A_α can be expressed in terms of the points B_α as follows

$$\begin{cases} A_0 = B_0 - pB_4, \\ A_1 = B_1 + 2pB_2 - p^2B_3 - 2pqB_0, \\ A_2 = B_2 - pB_3 - qB_0, \\ A_3 = B_3, \\ A_4 = B_4. \end{cases} \quad (14)$$

Consider a displacement of the straight lines $B_1 \wedge B_2$ along the hypersurface F . Suppose that Z is an arbitrary point of this straight line,

$$Z = B_1 + \lambda B_2. \quad (15)$$

Differentiating (15) and taking into account (14) and $dA_\alpha = 0$, we find that

$$dZ \equiv (2qdp + \lambda dq)B_0 + \lambda dp(B_3 + qB_4) \pmod{B_1, B_2}. \quad (16)$$

It follows from relation (16) that

1. A tangent hyperplane to the hypersurface F is spanned by the points B_1, B_2, B_0 and $B_3 + qB_4$. This hyperplane is fixed when the point Z moves along the straight line $B_1 \wedge B_2$. Thus the hypersurface F is tangentially degenerate of rank 2, and the straight lines $B_1 \wedge B_2$ form a Monge-Ampère foliation on F .
2. The system of equations

$$\begin{cases} 2qdp + \lambda dq &= 0, \\ \lambda dp &= 0 \end{cases} \quad (17)$$

defines singular points on the straight line $B_1 \wedge B_2$, and on the hypersurface F it defines torse. The system of equations (17) has a nontrivial solution with respect to dp and dq if and only if its determinant vanishes: $\lambda^2 = 0$. Hence by (15), a singular point on the straight line $B_1 \wedge B_2$ coincides with the point B_1 . For $\lambda = 0$, system (17) implies that $dp = 0$, i.e., $p = \text{const.}$ Thus it follows from (9) that the point $B_1 \in C$ is fixed, and as a result, the torse corresponding to this constant parameter p is a pencil of straight lines with the center B_1 located in the 2-plane $\tau(p) = B_0 \wedge B_1 \wedge B_2$.

3. All singular points of the hypersurface F belong to the conic $C \subset P^\infty$ defined by equation (8). Thus if we consider the hypersurface F in an Euclidean space R^4 , then on F there are no singular points in a proper part of this space.
4. The hypersurface F considered in the proper part of an Euclidean space is not a cylinder since its rectilinear generators do not belong to a bundle of parallel straight lines. A two-parameter family of rectilinear generators of F decomposes into a one-parameter family of plane pencils of parallel lines.

4. No one of properties 1–4 characterizes Bourgain’s hypersurfaces completely: they are necessary but not sufficient for these hypersurfaces. The following theorem gives a necessary and sufficient condition for a hypersurface to be of Bourgain’s type.

Theorem 1 *Let l be a proper straight line of an Euclidean space R^4 enlarged by the plane at infinity P_∞^3 , and let C be a conic in the 2-plane σ . Suppose that the straight line l and the conic C are in a projective correspondence. Let $B_0(p)$ and $B_1(p)$ be two corresponding points of l and C , and let τ be the 2-plane passing through the point B_0 and tangent to the conic C at the point B_1 . Then*

- (a) *when the point B_0 is moving along the straight line l , the plane τ describes a Bourgain hypersurface, and*
- (b) *any Bourgain hypersurface satisfies the above construction.*

Proof. The necessity (b) of the theorem hypotheses follows from our previous considerations. We prove the sufficiency (a) of these hypotheses. Take a fixed frame $\{A_u\}$, $u = 0, 1, 2, 3, 4$, in the space R^4 enlarged by the plane at infinity P_∞^3 as follows: its point A_0 belongs to l , the point A_4 is the point at infinity of l , and the points A_1, A_2 , and A_3 are located at the 2-plane at infinity σ in such a way that a parametric equation of the straight line l is $B_0 = A_0 + pA_4$, and the equation of C has the form (9). The plane τ is defined by the points B_0, B_1 , and $\frac{dB_1}{dp}$. The parametric equations of this plane have the form (12). Excluding the parameters α, β, γ , and p from these equations, we will return to the cubic equation (4) defining the Bourgain hypersurface B in homogeneous coordinates. ■

The method of construction of the Bourgain hypersurface used in the proof of Theorem 1 goes back to the classical methods of projective geometry developed by Steiner [St 32] and Reye [R 66].

5. In conclusion we prove the following theorem.

Theorem 2 *The Sacksteder hypersurface S and the Bourgain hypersurface B are locally equivalent, and the former is the standard covering of the latter.*

Proof. In an Euclidean space R^4 , in Cartesian coordinates x_1, x_2, x_3, x_4 , the equation of the Sacksteder hypersurface S (see [S 60]) has the form

$$x_4 = x_1 \cos x_3 + x_2 \sin x_3. \quad (18)$$

The right-hand side of this equation is a function on the manifold $M^3 = \mathbb{R}^2 \times S^1$ since the variable x_3 is cyclic. Equation (18) defines a hypersurface on the manifold $M^3 \times S^1$. The circumference $S^1 = \mathbb{R}/2\pi\mathbb{Z}$ has a natural projective structure of P^1 . In the homogeneous coordinates $x_3 = \frac{u}{v}$, the mapping $S^1 \rightarrow P^1$, can be written as $x^3 \rightarrow (u, v)$. By removing the point $\{v = 0\}$ from S^1 , we obtain a 1-to-1 correspondence

$$S^1 - \{v = 0\} \longleftrightarrow \mathbb{R}^1. \quad (19)$$

Now we can consider the Sacksteder hypersurface S in R^4 or, if we enlarge R^4 by the plane at infinity P_∞^3 , in the space P^4 .

Next we show how by applying the mapping $S^1 \rightarrow P^1$, we can transform equation (18) of the Sacksteder hypersurface S into equation (4) of the Bourgain hypersurface B . We write this mapping in the form

$$x_3 = 2 \arctan \frac{u}{v}, \quad \frac{u}{v} \in R, \quad |x_3| < \pi. \quad (20)$$

It follows from (20) that

$$\begin{cases} \frac{u}{v} = \tan \frac{x_3}{2}, \\ \cos x_3 = \frac{1 - \tan^2 \frac{x_3}{2}}{1 + \tan^2 \frac{x_3}{2}} = \frac{v^2 - u^2}{v^2 + u^2}, \\ \sin x_3 = \frac{2 \tan \frac{x_3}{2}}{1 + \tan^2 \frac{x_3}{2}} = \frac{2uv}{v^2 + u^2}. \end{cases} \quad (21)$$

Substituting these expressions into equation (18), we find that

$$x_4(u^2 + v^2) = x_1(v^2 - u^2) + 2x_2uv,$$

i.e.,

$$(x_4 + x_1)u^2 + (x_4 - x_1)v^2 - 2x_2uv = 0. \quad (22)$$

Make a change of variables

$$z_1 = x_4 - x_1, \quad z_2 = -2x_2, \quad z_3 = x_1 + x_4, \quad z_0 = u, \quad z_4 = v.$$

As a result, we reduce equation (22) to equation (4). It follows that the Sacksteder hypersurface S defined by equation (18) is locally equivalent to the Bourgain hypersurface defined by equation (4).

Note also that if the cyclic parameter x_3 changes on the entire real axis R , then we obtain the standard covering of the Bourgain hypersurface B by means of the Sacksteder hypersurface S . ■

References

- [A 87] Akivis, M. A., *On multidimensional strongly parabolic surfaces*, (Russian) Izv. Vyssh. Uchebn. Zaved. Mat. **1987**, no. 5 (311), 3–10. English translation: Soviet Math. (Iz. VUZ) **31** (1987), no. 5, 1–11.
- [AG 93] Akivis, M. A., and V. V. Goldberg, *Projective differential geometry of submanifolds*, North-Holland, Amsterdam, 1993, xii+362 pp.
- [AGL] Akivis, M. A., V. V. Goldberg, and J. Landsberg, *On the Griffiths-Harris conjecture on varieties with degenerate Gauss mappings*, 1999, 3 pp. (submitted).
- [C 39] Cartan, É., *Sur des familles remarquables d'hypersurfaces isoparamétriques dans les espaces sphériques*, Math. Z. **45**, 335–367; see also *Œuvres complètes*, Partie **III**, *Géométrie différentielle. Divers*, pp. 1447–1479, Editions du C. N. R. S., Paris, 1984.
- [FW 95] Fischer, G. and H. Wu, *Developable complex analytic submanifolds*, Internat. J. Math. **6** (1995), no. 2, 229–272.
- [GH 79] Griffiths, P. A. and J. Harris, *Algebraic geometry and local differential geometry*, Ann. Sci. École Norm. Sup. (4) **12** (1979), 355–452.
- [I 98] Ishikawa, G., *Developable hypersurfaces and algebraic homogeneous spaces in a real projective space*, in *Homogeneous structures and theory of submanifolds* (Kyoto, 1998). Sūrikaiseikikenkyūsho Kōkyūroku No. 1069 (1998), 92–104.
- [I 99a] Ishikawa, G., *Singularities of developable surfaces*, *Singularity theory (Liverpool, 1996)*, xxii–xxiii, 403–418, London Math. Soc. Lecture Note Ser., 263, Cambridge Univ. Press, Cambridge, 1999.

- [I 99b] Ishikawa, G., *Developable hypersurfaces and homogeneous spaces in a real projective space*, Lobachevskii J. Math. **3** (1999), 113–125.
- [IM 97] Ishikawa, G. and T. Morimoto, *Solution surfaces of Monge-Ampère equations*, Hokkaido Univ. Preprint Series **376** (1997), 16 pp.
- [L 99] Landsberg, J. M., *Algebraic geometry and projective differential geometry*, Lecture Notes Series, No. 45, Seoul National Univ., Seoul, Korea, 1999, 85pp.
- [M 94] H. Mori, *Remarks on complete deformable hypersurfaces in R^4* , J. Differential Geom. **40** (1994), no. 1, 1–6.
- [R 68] Reye, Th., *Geometrie der Lage*, 2nd ed., 2 vol., Rümpler, Hannover, 1868; English transl.: *Lectures on the geometry of positions*, The Macmillan Comp., New York, 1898.
- [S 60] Sacksteder, R: *On hypersurfaces with no negative sectional curvature*, Amer. J. Math. **82** (1960), no. 3, 609–630.
- [St 32] Steiner, J., *Systematische Entwicklung der Abhängigkeit geometrischer Gestalten von einander*, Teil I und II, Berlin, 1832; 2nd ed., W. Engelmann, Leipzig, 1896, 126 pp. & 162 pp.; see also in Steiner, J., *Gesammelte Werke*, B. I, G. Reimer, Berlin, 1881.
- [W 95] Wu, H., *Complete developable submanifolds in real and complex Euclidean spaces*, Internat. J. Math. **6** (1995), no. 3, 461–489.
- [WZ 99] Wu, H. and P. Zheng, *On complete developable submanifolds in complex Euclidean spaces*, Preprint, 30 pp. (July 15, 1999).

Authors' addresses:

M. A. Aklivis
 Department of Mathematics
 Jerusalem College of Technology—Mahon Lev
 Havaad Haleumi St., P. O. B. 16031
 Jerusalem 91160, Israel

E-mail address: akivis@avoda.jct.ac.il

V. V. Goldberg
 Department of Mathematical Sciences
 New Jersey Institute of Technology
 University Heights
 Newark, N.J. 07102, U.S.A.

E-mail address: vlgold@m.njit.edu